

**Featured Items**

Soma
Tetracycline
Adipex-P
Fioricet
Tramadol

Online Drugstore Prices

A - M N - Z

Health Directory

Weight Loss
Antibiotics
Pain Relief
Muscle Relaxers
Sleeping Aids
Anxiety
Women's Health
Blood Pressure
Sexual Health
Headache
Allergy Medicine
Anti-Depressant
Anti-Psychotics
Chest Pain
Herpes
Men's Health
Skin Care
Asthma Medicine
Diabetes
Cholesterol
Bladder Control
Quit Smoking
Upset Stomach
Motion Sickness

Buy Esgic Plus

Buy Esgic Plus from the online drugstore by choosing from the options below.

Product	Qty/ Tubes	Description	Price	
	30	500mg Tablets	\$97	Buy Esgic Plus
Esgic Plus	60	500mg	\$140	Buy Esgic Plus
	90	500mg	\$214	Buy Esgic Plus

Esgic Plus Information (Zebutal)**Uses**

Esgic Plus is an analgesic, barbiturate, and stimulant combination used to treat tension headaches.

How To Use

Follow the directions for using Esgic Plus provided by your doctor. STORE ESGIC PLUS at room temperature in a tightly-closed container, away from heat and light. IF YOU MISS A DOSE OF ESGIC PLUS and you are using it regularly, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

Side Effects

SIDE EFFECTS, that may go away during treatment, include drowsiness, dizziness, lightheadedness, or nausea. If they continue or are bothersome, check with your doctor. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist.

Precautions

DO NOT EXCEED THE RECOMMENDED DOSE or take Esgic Plus for longer than prescribed without checking with your doctor. Exceeding the recommended dose or taking Esgic Plus for longer than prescribed may be habit-forming. AVOID ALCOHOL while you are using Esgic Plus. Esgic Plus will add to the effects of alcohol and other depressants. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to Esgic Plus. Using Esgic Plus alone, with other medicines, or with alcohol may lessen your ability to drive or to perform other potentially dangerous tasks. ESGIC PLUS CONTAINS ACETAMINOPHEN. Do not take additional acetaminophen for pain or fever without checking with your doctor or pharmacist. Ask your pharmacist if you have questions about which medicines contain acetaminophen. Acetaminophen may cause liver damage. If you drink alcohol on a daily basis, do not take Esgic Plus without first discussing it with your doctor. Alcohol use combined with acetaminophen may increase your risk for liver damage. Pain medications work best if they are used as the first signs of pain (or first signs of migraine) occur. If you wait until the pain has significantly worsened, the pain medicine may not work as well. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. FOR WOMEN TAKING BIRTH CONTROL PILLS: Esgic Plus may decrease the effectiveness of your birth control pills. To prevent pregnancy, use an additional form of birth control while you are taking Esgic Plus. IF YOU PLAN ON BECOMING PREGNANT, discuss with your doctor the benefits and risks of using Esgic Plus

Female Health

Actonel
Diflucan
Esgic Plus
Evista
Famvir
Fosamax
Ortho Evra Patch
Ortho Tri Cyclen
Prempro
Triphasil
Yasmin
Zebutal

during pregnancy. **ESGIC PLUS IS EXCRETED IN BREAST MILK. DO NOT BREAST-FEED** while taking Esgic Plus.

Drug Interactions

Drug interactions can result in unwanted side effects or prevent a medicine from doing its job. Use our drug interaction checker to find out if your medicines interact with each other.

Overdose

If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include vomiting, excessive sweating, dark urine, stomach pain, extreme fatigue, unusual drowsiness or dizziness, lack of feeling alert, slow or shallow breathing, cold or clammy skin, and loss of consciousness.

Notes

IF YOUR HEADACHE SYMPTOMS do not improve or if they become worse, contact your doctor. **DO NOT SHARE ESGIC PLUS** with others for whom it was not prescribed. **DO NOT USE ESGIC PLUS** for other health conditions. **KEEP ESGIC PLUS** out of the reach of children. **IF USING ESGIC PLUS FOR AN EXTENDED PERIOD OF TIME**, obtain refills before your supply runs out.

[View Page as PDF](#)

Find a prescription drug by clicking on the first letter of the drug's name to show a list of all drugs starting with that letter:

A B C D E F G H I J K L M N O P Q R S T U V W X Y Z

Marketing Partners

