



Featured Items

Soma
Tetracycline
Adipex-P
Fioricet
Tramadol

Online Drugstore
Prices

A - M N - Z

Health Directory

Weight Loss
Antibiotics
Pain Relief
Muscle Relaxers
Sleeping Aids
Anxiety
Women's Health
Blood Pressure
Sexual Health
Headache
Allergy Medicine
Anti-Depressant
Anti-Psychotics
Chest Pain
Herpes
Men's Health
Skin Care
Asthma Medicine
Diabetes
Cholesterol
Bladder Control
Quit Smoking
Upset Stomach
Motion Sickness

Buy Fioricet

Buy Fioricet from the online drugstore by choosing from the options below.

Product	Qty	Description	Price	
	30	Tablets	\$94.30	Buy Fioricet
Fioricet 50/325/40	60	Tablets	\$138.10	Buy Fioricet
	90	Tablets	\$189	Buy Fioricet
	30	Tablets	\$79.25	Buy Fioricet
Generic Fioricet 50/325/40	60	Tablets	\$89.20	Buy Fioricet
(Butalbital, Apap, Caffeine)	90	Tablets	\$89	Buy Fioricet

Fioricet Information

IMPORTANT NOTE: The following information is intended to supplement, not substitute for, the expertise and judgment of your physician, pharmacist or other healthcare professional. It should not be construed to indicate that use of the drug is safe, appropriate, or effective for you. Consult your healthcare professional before using this drug.

BUTALBITAL/ACETAMINOPHEN/CAFFEINE - ORAL (byou-TAL-bih-tall/uh-seet-uh-MEE-no-fen/kaff-EEN)

COMMON BRAND NAME(S): Esgic, Fioricet

USES: This medication is a pain reliever and sedative. It is used to relieve mild to moderate pain and tension headaches.

HOW TO USE: To prevent upset stomach, take this medication with food or milk. Pain medications work best in preventing pain before it occurs. Once the pain becomes intense, the medication is not as effective in relieving it. Take this as prescribed. Do not increase your dose or take it more frequently without your doctor's approval.

SIDE EFFECTS: May cause lightheadedness, dizziness, drowsiness, stomach upset, nausea, and sleeping disturbances the first few days as your body adjusts to the medication. If these symptoms persist or become bothersome, inform your doctor. Notify your doctor immediately if you develop: a rapid/irregular heartbeat. An allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include: rash, itching, swelling, severe dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS: Tell your doctor your medical history, especially of: serious liver disease, alcohol or narcotic dependence, emotional/mental conditions, heart disease (arrhythmias, recent MI), stomach/intestinal ulcers, any allergies. Acetaminophen may cause liver damage. Daily use of alcohol, especially when combined with acetaminophen, may increase your risk for liver damage. Check with your doctor or pharmacist for more information. To prevent oversedation, avoid using alcohol and other sedative type medications while taking this. This

Pain Relief

Bextra
Celebrex
Diclofenac
Fioricet
Naproxen
Tramadol
Ultracet
Ultram
Vioxx

drug may make you dizzy or drowsy; use caution engaging in activities requiring alertness such as driving or using machinery. Avoid alcoholic beverages. This medication contains caffeine. Avoid drinking large amounts of caffeinated beverages while taking this medication as excessive nervousness and irritability can occur. This medication should be used during pregnancy only if clearly needed. Small amounts of this medication appear in breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS: Tell your doctor of all prescription and nonprescription drugs you may use, especially of: "blood thinners" Tell your doctor or pharmacist of all prescription and nonprescription drugs you may use that cause drowsiness such as: medicine for sleep (e.g., sedatives), tranquilizers, anti-anxiety drugs (e.g., diazepam), narcotic pain relievers (e.g., codeine), psychiatric medicines (e.g., phenothiazines such as chlorpromazine, or tricyclics such as amitriptyline), anti-seizure drugs (e.g., carbamazepine), muscle relaxants, antihistamines that cause drowsiness (e.g., diphenhydramine). Because this medication contains acetaminophen, check with your doctor or pharmacist before taking other medications containing acetaminophen which is commonly found in nonprescription pain relievers and cough-and-cold products. Read labels carefully to check ingredients. This drug may interfere with the effectiveness of birth control pills. Discuss using other methods of birth control with your doctor. Do not start or stop any medicine without doctor or pharmacist approval.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include vomiting, unusual drowsiness, lack of feeling alert, slow or shallow breathing, cold or clammy skin, loss of consciousness, dark urine, stomach pain, and extreme fatigue.

NOTES: Do not share this product with others.

MISSED DOSE: If you miss a dose, take it as soon as remembered; do not take if it is almost time for the next dose, instead, skip the missed dose and resume your usual dosing schedule. Do not "double-up" the dose to catch up.

STORAGE: Store at room temperature away from moisture and sunlight. Do not store in the bathroom.

[View page as PDF](#)

Find a prescription drug by clicking on the first letter of the drug's name to show a list of all drugs starting with that letter:

A B C D E F G H I J K L M N O P Q R S T U V W X Y Z

Marketing Partners

